



SRI KONDA LAXMAN TELANGANA HORTICULTURAL UNIVERSITY
Administrative office, Mulugu (V&M), Siddipet District- Telangana

Weather Based Agromet Advisories for Telangana in Horticultural Crops
01-10-2026 to 07-10-2026

Weather during past Three Days

During the last three days, generally sunny weather prevailed across Telangana. Maximum temperatures were around 35°C, while minimum temperatures ranged from 22–23°C.

Weather Forecast for the Next Five Days

During the next five days, partly cloudy weather is expected across Telangana. Light rainfall accompanied by thunderstorms is likely at isolated places on October 3. On the remaining days, rainfall is expected to be low, with sunny weather likely to prevail.

- Average maximum temperature: 34°C–36°C
- Average minimum temperature: 22°C–26°C

Day 1 (From: 01/10/2026 08:30 hrs to 02/10/2026 08:30 hrs)

Dry Weather very likely to prevail over Telangana.

Day 2 (From: 02/10/2026 08:30 hrs to 03/10/2026 08:30 hrs)

Light to Moderate Rain or Thundershowers very likely to occur at isolated places over Telangana. Thunderstorm accompanied with Lightning and Gusty Winds (30-40 kmph) very likely to occur at isolated places in Jayashankar Bhupalpally, Mulugu, Bhadrachalam, Kothagudem, Khammam, Nalgonda, Suryapet, Nagarkurnool districts of Telangana

Day 3 (From: 03/10/2026 08:30 hrs to 04/10/2026 08:30 hrs)

Light to Moderate Rain or Thundershowers very likely to occur at isolated places over Telangana. Thunderstorm accompanied with Lightning and Gusty Winds (30-40 kmph) very likely to occur at isolated places in Mulugu, Bhadrachalam, Kothagudem, Khammam, Nalgonda, Suryapet, Mahabubnagar, Nagarkurnool, Wanaparthy, Narayanpet, Jogulamba Gadwal districts of Telangana

Day 4 (From: 04/10/2026 08:30 hrs to 05/10/2026 08:30 hrs)

Dry Weather very likely to prevail over Telangana.

Day 5 (From: 05/10/2026 08:30 hrs to 06/10/2026 08:30 hr)

Dry Weather very likely to prevail over Telangana.

VEGETABLES

- In view of the expected high temperature during the next five days, flowering and fruit-setting stages of vegetable crops may be affected. Therefore, irrigation should be provided to maintain adequate soil moisture. Farmers in water-scarcity areas are advised to adopt drip and sprinkler irrigation methods instead of conventional irrigation practices.

- The predicted weather conditions may also favour the incidence of sucking pests such as whiteflies and thrips. For their monitoring and management, blue and yellow sticky traps should be installed in the Field and monitored regularly. If pest incidence exceeds the economic threshold level (ETL), Imidacloprid 17.8% SL @ 0.5 mL per litre of water may be sprayed.
- **Chilli**
 - In view of the weather forecast, farmers are advised to monitor black thrips, whiteflies and mites regularly.
 - Install blue sticky traps @ 8–10/acre for thrips and yellow sticky traps @ 8–10/acre for whiteflies.
 - If incidence is high, spray Spinosad 45% SC @ 0.3 ml/L for thrips and Spinetoram 11.7% SC @ 0.5 ml/L for whiteflies.
 - Under dry and warm conditions, monitor young leaves for mites and, if necessary, spray Spiromesifen 22.9% SC @1 ml/L or Abamectin 1.9% EC @ 0.5 ml/L.

Fruits

- **Mango**
 - Dislodge the nests of leaf-webber using bamboo poles and spray Chlorpyrifos (2 ml/L) or Acephate (1.5 g/L).
 - To managehoppershibernating in the tree bark, spray Acephate @ 1.5 g/L or Imidacloprid @ 0.3 ml/L of water by thoroughly drenching the tree.
- **Guava**
 - Collect and destroy the canker affected fruits, dried twigs and spray with lambda cyhalothrin @ 0.5-1.0 ml/lit + (Carbendazim + Mancozeb) @ 2g/L for control of damage by Tea mosquito bug.Avoid pesticide spraying during rains.
 - To monitor fruit fly in guava, install methyl eugenol pheromone traps @ 4-8/acre.
- **Acid lime and Sweet orange** - In view of scanty rainfall or showers forecast,after cessation of rains, spray Copper Oxychloride 50 WP @ 3 g/litre of water in Citrus orchards as prophylactic measure to control the citrus canker.

PLANTATION, SPICES, MEDICINAL AND AROMATIC PLANTS

- **Turmeric**
 - As a dry spell is expected to persist throughout the week, irrigation should be provided as needed. 2-3 irrigations for heavy soils and 3-5 irrigations to light soils can be given.
 - To control shoot borer incidence chlorantraniliprole@0.5ml/lit or flubendiamide @0.3 ml/lit can be given as a foliar spray.
- **Ginger** - To control leaf roller incidence chlorantraniliprole @0.5ml/lit can be given as a foliar spray
- **Oil palm** - Regularly monitor the palms for bud rot and bunch rot symptoms. If symptoms are observed, spray Propiconazole + Difenconazole @ 1 ml/L or Carbendazim + Mancozeb @ 3 g/L, on crown region and bunches.
- **FLOWERS** - Apply water-soluble fertilizers directly through the drip system to ensure uniform nutrient distribution and immediate uptake.